

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>A61K 38/30</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 97/21449</b> (43) International Publication Date: 19 June 1997 (19.06.97)</p>
<p>(21) International Application Number: PCT/US96/19663 (22) International Filing Date: 11 December 1996 (11.12.96) (30) Priority Data: 08/571,802 13 December 1995 (13.12.95) US (71) Applicant: COLORADO STATE UNIVERSITY RESEARCH FOUNDATION [US/US]; 601 South Howes, Fort Collins, CO 80523 (US). (72) Inventor: ISHII, Douglas, N.; 3301 Rawhide Drive, La Porte, CO 80535 (US). (74) Agent: WAACK, Janelle, D.; Arnold, White &amp; Durkee, P.O. Box 4433, Houston, TX 77210 (US).</p>		<p>(81) Designated States: CA, CN, JP, PL, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE)  Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: METHOD FOR EFFECTING CHANGES IN THE CENTRAL NERVOUS SYSTEM BY ADMINISTRATION OF IGF-I OR IGF-II (57) Abstract  The invention concerns a method for effecting changes in the central nervous system, including treatment of brain and spinal cord disorders or diseases, by parenteral administration of insulin-like growth factors I (IGF-I) or insulin-like growth factor II (IGF-II).</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHOD FOR EFFECTING CHANGES  
IN THE CENTRAL NERVOUS SYSTEM  
BY ADMINISTRATION OF IGF-I OR IGF-II

5

The Government may own certain rights in the present invention pursuant to National Institute of Neurological Disorders and Stroke Grant No. 5R01 NS24327 to Douglas N. Ishii.

10 The invention concerns a method for effecting changes in the central nervous system by administration of insulin-like growth factor I (IGF-I) or insulin-like growth factor II (IGF-II). More particularly, the invention is directed to a method for treating disorders or diseases of the brain or spinal cord by parenteral administration of IGF-I or IGF-II.

15 Many people suffer from disorders and diseases of the brain such as Alzheimer's Disease, Parkinson's Disease, dementia associated with Acquired Immunodeficiency Syndrome (AIDS), Pick's Disease, Huntington's Disease, memory loss due to aging, stroke, derangements of the intellect and behavior, neurologic effects of aging, and the like. The treatment of a brain disorder or disease is generally more complicated than treatment of the peripheral nervous  
20 system due to the blood-brain barrier which poses an additional obstacle to the delivery of pharmaceutical agents to the brain.

The endothelial cells lining the brain vasculature separate the brain from the blood. This "blood-brain barrier" has been reviewed in Friedemann (1942); Rowland et al. (1991); and Schlosshauer (1993). The blood-brain  
25 barrier protects the brain, e.g., from changes in circulating levels of ions, neurotransmitters and growth-altering factors. For example, if high concentrations of certain neurotransmitters were to enter the brain from blood,

brain neurons may become inappropriately activated and overexcitation might cause brain disorder or damage.

5 The brain capillaries which make up the blood-brain barrier are lined with endothelial cells cemented together with very tight junctions, which have few transendothelial channels and allow only scanty pinocytosis. By contrast, the capillaries of the peripheral tissues are lined with endothelial cells which are loosely cemented together with 30-80 Angstrom diameter pores at their junctions, and which have many more transendothelial channels and allow abundant pinocytosis. The tight junctions between the endothelial cells in the  
10 blood-brain barrier limit the kinds of molecules that can effectively cross the blood-brain barrier to enter the brain. These molecules include essential molecules needed for brain metabolism and for which there is a specific transport system, such as glucose and amino acids. In addition, small lipophilic molecules can dissolve in the lipoid environment of the endothelial cell plasma membrane and passively diffuse into the brain. By contrast, polar,  
15 ionized and large molecules, including proteins, are typically excluded from the brain by the blood-brain barrier.

Similarly, the spinal cord is protected by a blood-spinal cord barrier. For example, spinal cord interneurons have their cell bodies and neuritic  
20 processes entirely within the blood-spinal cord barrier. As for the brain, there is a need for a method to effect changes in or treat the mature spinal cord, particularly following traumatic spinal cord injury.

Various procedures have been contemplated in efforts to circumvent the blood-brain barrier and effect changes in the brain. For example, in one  
25 approach, a small hole is drilled through the skull through which neurotrophic growth factors might be applied to the ventricles of the brain via a catheter, or through which injections or implants can be made. Implanted gel foam, tissues or cells might be used to release such growth factors into the brain.

However, such invasive procedures are understandably difficult, risky and require costly surgical procedure. Alternatively, it may be possible to encapsulate neurotrophic proteins within lipid vesicles, and use such vesicles to enhance delivery of factors across the blood-brain barrier.

5 In another approach, U.S. Patent No. 4,801,575 to Pardridge discloses chimeric peptides wherein a hydrophilic neuropeptide is conjugated via a covalent bond to a transportable peptide for delivering the neuropeptide to the brain. Pardridge discloses such chimeric peptides where the transportable peptide is insulin, transferrin, insulin-like growth factor I (IGF-I), insulin-like  
10 growth factor II (IGF-II), basic albumin or prolactin, and where the neuropharmaceutical agent is somatostatin, thyrotropin releasing hormone, vasopressin, alpha interferon or endorphin. However, Pardridge does not employ IGF-I or IGF-II as an agent which itself is effective for treating a brain disorder or disease.

15 With regard to the spinal cord, previous studies have shown that IGF-I and IGF-II can enhance neurite outgrowth in cultured embryonic rat spinal cord neurons (Ishii et al. (1989)). However, the use of IGF-I and IGF-II to treat neurons within the mature spinal cord, e.g. having a developed blood-spinal cord barrier, particularly for the treatment of disorders and diseases of  
20 the mature spinal cord, has not been investigated.

IGF-I and IGF-II are approximately  $M_r$  7500-7700 in size. The neurotrophic properties of the IGFs are discussed in Ishii and Recio-Pinto (1987), which is incorporated herein by reference. IGF receptors are found in brain tissues (Sara et al. (1982); Goodyear et al. (1984)) and are present on  
25 neurons and neuroglia cells. IGFs have been shown to prevent the death of cultured embryonic chick sensory and sympathetic neurons and to promote neurite outgrowth (Recio-Pinto et al. (1986)).

Researchers have reported that  $^{125}\text{I}$ -labeled IGF-I and IGF-II cross the blood-brain barrier and selectively accumulate in specific hypothalamic and anterior thalamic nuclei (Reinhardt and Bondy (1994)). However, those researchers acknowledged that the physiological consequence of IGFs' apparent ability to cross the blood-brain barrier required further study.

The effect of parenterally administered IGF-I or IGF-II on the central nervous system, particularly for treatment of brain and spinal cord disorders or diseases, has not yet been determined. Moreover, such determinations are not predictable in view of the blood-brain barrier and blood-spinal cord barrier as obstacles to drug delivery, particularly for large protein molecules such as IGF-I or IGF-II.

Therefore, there is a need in the art of biotechnology and the biopharmaceutical industries for a method of effecting changes to the central nervous system by the administration of large protein molecules across the blood-brain barrier and blood-spinal cord barrier, particularly where the method of treatment involves administration of IGF-I or IGF-II.

The invention concerns a method for effecting changes in the central nervous system by administration of IGF-I or IGF-II. In a preferred embodiment, the invention is directed to a method for treating disorders or diseases of the brain, such as Alzheimer's Disease, Parkinson's Disease, and AIDS-related dementia, by parenteral administration of IGF-I or IGF-II. In another preferred embodiment, the invention is directed to a method for treating disorders of or trauma to the spinal cord by parenteral administration of IGF-I or IGF-II. This invention relates to the use of IGFs to ameliorate damage to or treat brain and spinal cord disorders, disease, stroke or trauma by the use of parenteral administration of IGF-I and/or IGF-II in a pharmaceutical composition.

For purposes of this invention, several terms are defined below.

"IGF-I" refers to insulin-like growth factor I, and is also known as IGF1, IGF1, IGF-1 or somatomedin C. For purposes of the invention IGF-I also encompasses homologs of IGF-I from various animal species, whether extracted from tissues or derived as products of recombinant genetic expression vectors, and IGF molecules with substantial sequence homology to human and animal IGF-I that bind to type I IGF receptors. For purposes of the invention, IGF-I does not, however, encompass fusion proteins of IGF-I and a non-IGF peptide, wherein IGF-I functions as a transportable peptide and not as a pharmaceutical agent for treatment of a disorder or disease.

"IGF-II" refers to insulin-like growth factor II, and is also known as IGFII, IGF2 or IGF-2. The invention also encompasses homologs of IGF-II from various animal species, whether extracted from tissues or derived as products of recombinant genetic expression vectors, and IGF molecules with substantial sequence homology to human and animal IGF-II that bind to type I or type II IGF receptors. For purposes of the invention, IGF-II does not, however, encompass fusion proteins of IGF-II and a non-IGF protein, wherein IGF-II functions as a transportable peptide and not as a pharmaceutical agent for treatment of a disorder or disease.

"Brain" is defined as the mature (post-birth) brain within the blood-brain barrier. For purposes of this invention, the brain does not include circumventricular organs such as the pituitary.

"Spinal cord" is defined as the mature (post-birth) spinal cord contained within the blood-spinal cord barrier. For purposes of this invention, the spinal cord does not include neurons, such as motor neurons, whose axons lie in peripheral nerves outside of the blood-spinal cord barrier.

FIG. 1 is a graphical representation of the results of parenteral administration of IGF-I to diabetic and nondiabetic adult rats. The graph shows that administration of IGF-I effectively prevents impairment of IGF-II gene expression in the brain of adult diabetic rats.

5 FIG. 2A is a graphical representation of the limb withdrawal reflex (lower tracing) and muscle EMG (upper tracing) in a control rat exposed to surgery, but with no harm to the locus ceruleus cells.

FIG. 2B is a graphical representation of the limb withdrawal reflex (lower tracing) and muscle EMG (upper tracing) in the lesioned rat exposed to surgery, with damaged locus ceruleus cells and having a subcutaneously  
10 implanted miniosmotic pump releasing vehicle (no drug delivery).

FIG. 2C is a graphical representation of the limb withdrawal reflex (lower tracing) and muscle EMG (upper tracing) in the lesioned rat exposed to surgery, with damaged locus ceruleus cells and implanted subcutaneously with  
15 a miniosmotic pump releasing 4.8  $\mu$ g/day recombinant human IGF-II dissolved in a vehicle.

The invention concerns a method for effecting changes in the central nervous system by the parenteral administration of IGF-I or IGF-II. In a preferred embodiment, the invention is directed to the parenteral  
20 administration of IGF-I and/or IGF-II to effect a change in the brain or spinal cord, particularly where that change is a treatment for a brain or spinal cord disorder or disease.

Procedures to purify and obtain physiologically active IGF-I and IGF-II are known in the art (Zumstein and Humbel (1985); Svoboda and Van Wyk (1985)). IGF-I and IGF-II are commercially available as human recombinant  
25 factors, and are sold by Upstate Biotechnology, Inc., Lake Placid, New York; GroPep Ltd., Adelaide, Australia; Austral Biologicals, San Ramon, Calif.; and others.



Pharmaceutical compositions to be employed in the methods according to the invention comprise an effective amount of IGF-I or IGF-II. In a preferred embodiment, IGF-I or IGF-II is present in the pharmaceutical composition in an amount sufficient to have a therapeutic effect on the condition to be treated. In a further preferred embodiment, IGF-I or IGF-II is present in an amount from about 0.1% to 100% of the pharmaceutical composition. For example, the IGF-I or IGF-II may be administered in an amount from about 0.1 µg/kg/day up to about 4 mg/kg/day. As another example, the IGF-I or IGF-II may be administered in an amount from about 400 ng/kg/hour up to about 160 µg/kg/hour.

As one of ordinary skill in the art will appreciate, the dosage of pharmaceutical compositions can be adjusted as needed for a particular route of administration, weight of subject, and general condition and disorder or disease of the patient to be treated whether human or veterinary. Appropriate serum glucose monitoring should be done to prevent hypoglycemia, particularly when the higher end of the IGF dosage range is elected. The half-times of elimination, volumes of distribution, daily production rates, and serum concentrations are established pharmacokinetic parameters for IGF-I and IGF-II in normal humans (Guler et al. (1989); Zapf et al. (1981)).

The pharmaceutical compositions employed in the methods according to the invention may further comprise an inorganic or organic, solid or liquid, pharmaceutically acceptable carrier which is preferably suitable for parenteral administration. Compositions may optionally contain adjuncts including preservatives, wetting agents, emulsifiers, solubilizing agents, stabilizing agents, buffers, solvents and salts to maintain tonicity and osmotic pressure. Compositions may be sterilized and exist as solids, particulates or powders, solutions, suspensions or emulsions.

The methods according to the invention preferably employ parenteral administration. Any means of parenteral IGF administration may be utilized in this invention, including intradermal, subcutaneous, intramuscular, intravenous, intra-arterial or intraperitoneal administration. Other means for parenteral IGF administration might include use of a special infusion or slow release device, release of IGFs from implanted cells or device containing cells, or gene therapy into tissues, all outside the blood-brain barrier or blood-spinal cord barrier with the intent of effecting a change on the central nervous system, particularly to treat a brain or spinal cord disorder or disease.

The following examples are provided to enable those of ordinary skill in the art to make and use the methods of the invention. These examples are not intended to limit the scope of what the inventor regards as his invention. Efforts have been made to ensure accuracy with respect to numbers used to characterize the measured conditions; however, some experimental errors and deviations may be present.

## EXAMPLE I

### Example of a Pharmaceutical Preparation with IGF-I or IGF-II

A dry ampule (1-60 ml) is partially filled with a sterile solution of IGF-I or IGF-II, and optionally pharmaceutical adjuncts or carriers, and lyophilized. The parenteral solution is prepared by adding an appropriate volume of sterile water, saline or 0.001-0.1 M acetic acid. A pharmaceutical packet may contain a desired number of ampules for a course of treatment, optionally together with instructions for use.

## EXAMPLE II

### Parenteral Administration of IGF-I Having Effect on Impaired IGF-II Gene Expression in Metabolically Disordered Brain

5

Certain conditions may involve significant damage to the brain with a potential breakdown in the blood-brain barrier, for example, severe concussion injury to the brain, large penetrating wounds or infections causing inflammation to the meninges. The parenteral administration of IGFs under such conditions would not conclusively reveal whether IGFs can generally cross the blood-brain barrier in effective amounts and were therefore avoided in these examples.

10

This example involves the metabolic disorder of diabetes using the well-studied diabetic rat as a model. The blood-brain barrier is more permeable for certain small ions such as sodium and potassium, but not to other small ions and molecules such as chlorine, calcium or sucrose, in rats where diabetes is experimentally induced with streptozotocin (Knudsen et al. (1986); Jakobsen et al. (1987)). It is expected that, if chlorine, calcium or sucrose permeability is not increased in subjects having a diabetic condition, the permeability of the blood-brain barrier is not expected to be reduced for larger protein molecules, such as IGFs.

15

20

Adult rats (12-14 weeks old, about 300 grams in body weight) were treated with streptozotocin to induce insulin-deficient diabetes. One week later, diabetic rats were implanted subcutaneously in the mid-back with miniosmotic pumps releasing either recombinant human IGF-I (4.8 µg per day) or vehicle. After two weeks of continuous parenteral administration of IGF-I, the rat brains were stripped of the meninges and assayed for IGF-II

25

mRNA content. Standard biochemical techniques were used to extract RNA from brain and run Northern and slot blots. A rat cDNA containing the entire coding region of rat IGF-II was used to prepare a single-stranded anti-sense <sup>32</sup>P-labeled hybridization probe, which previously was well characterized. The probe does not cross-hybridize to IGF-I mRNA. Autoradiograms were used to measure IGF-II mRNA content in relative densitometric units (rel. units) per mg wet weight brain tissue. Values are means  $\pm$  SEM (N = 4 rats per group). The experimental methods are more completely described elsewhere (Soares et al. (1985); Ishii et al. (1994)).

As shown in FIG. 1, IGF-II mRNA content per mg wet weight tissue was significantly reduced in the brains of diabetic adult rats compared to that of nondiabetic rats. This defines a biochemical abnormality in diabetic brain. IGF-II mRNA is produced in various brain regions including hippocampus, thalamus, cerebral cortical layer 5, and choroid plexus (Hynes et al. (1988); Lee et al. (1992)). Neuroglia cells produce IGF-II mRNAs (Rotwein et al. (1988)). The inventor has also found IGF-II mRNA in brain regions such as hippocampus, striatum, midbrain, cerebellum and pons.

As further shown in FIG. 1, the impaired IGF-II gene expression in diabetic brain was ameliorated by parenteral IGF-I. These data show that parenteral administration of IGF-I effects a change in brain, and that it can correct a brain disorder resulting from a disease, in this case diabetes.

This correction of a brain disorder was not due to a significant reduction in hyperglycemia in diabetes, because these same low doses of IGF-I had no effect on hyperglycemia, although they did spare impaired sensory nerve regeneration in diabetic mature rats (Ishii and Lupien (1995)). Other studies have found that the same low doses of IGF-I or IGF-II have no significant effect on either hyperglycemia or weight loss in diabetic adult rats while hyperalgesia (pain) was prevented (Zhuang et al. (1994)). At many-fold

higher IGF doses, hyperglycemia or weight loss can be reduced, but this is not required for IGFs to act on the nervous system.

### EXAMPLE III

5

#### **Parenteral Administration of IGF-II Having Effect on an Impaired Behavior Resulting from a Brain Lesion**

10

In this model, a chemical lesion produced death of locus ceruleus cells in adult rat brain. Locus ceruleus cells normally project their noradrenergic axons down into the spinal cord to synapse on interneurons which, in turn, modify the activity of motor neurons controlling the hind limb withdrawal reflex. This example tested the capacity of parenterally administered recombinant human IGF-II to preserve this withdrawal reflex. Such preservation would result only if the administered IGF-II were to act on the brain to preserve the noradrenergic axons.

15

20

25

In the experiment shown in FIG. 2, 6-hydroxydopamine (6OHDA) (4  $\mu$ l of a solution consisting of 12.5 mg 6OHDA per ml and 0.2 mg/ml ascorbic acid in isotonic saline) was injected via a 30 gauge needle into the cisterna magna of adult (12-week-old) Sprague Dawley rats to destroy the noradrenergic locus ceruleus cells in the pons of the brain. Such treatment resulted in the disappearance of the axons from these noradrenergic cells which normally descend down the spinal cord to synapse on interneurons, which, in turn, modulated the activity of motor neurons controlling the limb withdrawal reflex. This reflex was measured by administration of 6.25 mg L-DOPA followed by electrical stimulation which caused release of noradrenaline from the descending noradrenergic fibers of the spinal cord.

leading to a large amplitude and long-latency hind limb withdrawal reflex whose force was monitored with a force-displacement transducer. The EMG in muscle was also recorded. Additional experimental details are available from Barnes et al. (1989) and Pulford et al. (1994).

5           FIG. 2A shows a control rat exposed to surgery and injected with solvent not containing 6OHDA. Under this condition there would be no harm to the locus ceruleus cells. The lower tracings show the intact large amplitude and long-latency hind limb withdrawal reflex force. F1-F4 are the force-time tracings which resulted from electrical stimulations of increasing intensity from 1.0-7.5 mA in 2.5 mA increments. The upper tracing shows the  
10           associated highly active EMG.

          FIG. 2B shows rats lesioned with 6OHDA that were implanted subcutaneously with miniosmotic pumps releasing vehicle alone. The lower tracings show loss of the duration and amplitude of the hind limb reflex force.  
15           The upper tracing shows loss of EMG activity.

          FIG. 2C shows rats lesioned with 6OHDA that were implanted subcutaneously with miniosmotic pumps releasing 4.8 µg/day recombinant human IGF-II. This treatment spared both the hind limb reflex and the EMG activity.

20           Treatment with 6OHDA is well established by others to result in the loss of noradrenergic brain cells. This was evidenced by the associated loss of the hind limb withdrawal reflex that was measured two weeks later in adult rats (FIG. 2A vs. 2B). A typical example is shown in FIG. 2B. The lower tracing shows the response to the L-DOPA test wherein the large amplitude and long-latency withdrawal reflex present in control unlesioned animals was  
25           almost entirely lost in the lesioned rat. The upper tracing shows the loss of EMG activity. These animals were implanted with subcutaneous osmotic minipumps that released vehicle only. In a group of six rats so treated, the

average peak force generated was  $15.6 \pm 3.8$  grams (mean, SD) in the hind limb withdrawal test.

By contrast, the hind limb withdrawal reflex was spared in 6OHDA lesioned animals implanted with subcutaneous pumps that released IGF-II (4.8  $\mu$ g/rat/day) continuously for two weeks. A typical example is shown in FIG. 2C. The lower tracing shows retention of the large amplitude and long-latency withdrawal reflex typical of unlesioned rats. The upper tracing shows retention of EMG activity. In a group of seven rats so treated, the average peak force was  $298 \pm 38$  grams (mean, SD). Therefore, IGF-II treatment caused a significant ( $P < 0.025$ ) sparing of the reflex vs. vehicle treatment. Thus, the parenteral administration of IGF ameliorated damage, most likely to the noradrenergic locus ceruleus nerve cells in the brain.

The reflex is mediated by noradrenaline, because the response to the L-DOPA test could be blocked by the  $\alpha$ -adrenergic blocking agent phentolamine.

The procedure producing lesions in the brain does not directly affect the spinal cord motoneurons, because direct electrical stimulation of the motor neuron axons in lesioned animals can cause effective contraction of the hind limb.

During the course of treatments in the above experiments, there were no indications of toxicity due to IGF administration.

The experimental results herein show the new and unexpected observation that parenterally administered IGF-I and IGF-II can effect changes in mature brains of treated rats. Parenteral administration can now be considered a viable route of administration for IGFs for the treatment of brain disorders and diseases, such as Alzheimer's Disease, Parkinson's Disease, dementia associated with Acquired Immunodeficiency Syndrome (AIDS), Pick's Disease, Huntington's Disease, memory losses due to aging, stroke, derangements of the intellect and behavior, neurologic effects of aging, and

the like. IGFs may also be useful for treatment of traumatic or chemical injury to the brain.

The abnormalities in human diabetic brain are well recognized (reviewed by Mooradian et al. (1988); McCall (1991)). These include major depression and cognitive deficits such as loss of memory and complex reasoning skills. Brain atrophy with degeneration of neurons, as well as axonal loss and neurotransmitter imbalances are observed. These are seen in diabetic humans (Reske-Nielsen et al. (1965); Olsson et al. (1968); Soininen et al. (1992)) as well as diabetic rats (Jakobsen et al. (1987); Trulsson et al. (1986)).

IGF-II is the predominant IGF in brain, and the decline in IGF-II mRNA content in diabetic brain (FIG. 1) suggests that it contributes to the pathogenesis of encephalopathy because IGFs are believed to be necessary for neuron survival, neurite outgrowth and maintenance of synapses. The parenteral administration of IGF-I prevented the decline in IGF-II mRNA within the context of diabetes, and it is expected that such parental administration will be useful to prevent or treat various biochemical disturbances in the diseased, disordered or injured brain.

It is believed that the decline in circulating IGF activity in diabetic patients contributes to abnormalities in diabetic brain. Inasmuch as liver is the primary source of circulating IGFs, other conditions that reduce liver function may contribute to associated brain disorders. For example, in hepatic encephalopathy chronic liver disease or liver failure is associated with altered behavior, impaired cognition, confusion, and coma. The latter carries risk of death or permanent neurological disability. Therefore, the invention further applies to the treatment of the brain in hepatic encephalopathy.

The locus ceruleus cells are located below the cerebellum deep within the pons in the rostral pontine central gray region. Their axons have terminals on interneurons located entirely within the spinal cord. Because these locus



ceruleus noradrenergic neurons and their axons are located entirely within the central nervous system enveloped completely by the blood-brain and blood-spinal cord barriers, it is clear that the parenteral administration of IGF-II effected a change in brain and/or spinal cord (FIG. 2). The parenteral administration of IGF-II was able to prevent the consequences of brain injury.

Other studies showed that a single treatment with 2  $\mu$ g IGF-II mixed together with the 6OHDA (injected through the cisterna magna) did not spare the hind limb withdrawal reflex. Thus, the continuous administration of IGF-II for two weeks was more effective.

IGF-II may have also prevented the secondary consequences of acute injury. For example, following an injury to brain, there can be secondary death of neurons over several days. Because the continuous administration of IGF-II for two weeks was effective whereas a single injection to the brain was not, the continuous administration of IGF appears to provide prophylaxis against secondary death of neurons or functional damage to neurons.

Although the most likely explanation is that IGFs crossed the blood-brain barrier in effective amounts, this invention encompasses methods in which IGFs are administered to effect changes on the brain, either directly or indirectly.

The methods of the invention are expected to be suitable for the treatment of other brain and spinal cord disorders and diseases such as Parkinson's Disease and Alzheimer's Disease. In Parkinson's Disease, there is a loss of dopaminergic neurons in the brain, as well as the noradrenergic neurons in the locus ceruleus. The invention shows that parenteral administration of IGFs can effect the brain system involving these noradrenergic neurons, and such treatment with IGFs is expected to help treat Parkinson's Disease. The dopaminergic and noradrenergic neurons are examples of closely related catecholaminergic neurons, and brain

catecholaminergic neurons are lost in Alzheimer's Disease. Neurofibrillary tangles are also observed in the locus ceruleus in Alzheimer's Disease, and such tangles are considered pathogenic. IGFs are expected to help support such locus ceruleus cells, catecholaminergic neurons, and other brain cells weakened in this disease.

The actions of IGFs are not confined to catecholaminergic and locus ceruleus brain cells, and intracranial IGF administration can support the survival of a large variety of different brain cells following ischemic injury. Thus, IGF parenteral treatment is expected to be useful in other disorders as well, including stroke, lobar atrophy (Pick's disease), Huntington chorea, and various neurodegenerative disorders where many types of neurons are at risk.

Many environmental neurotoxins are well known to cause damage to the brain, for instance, 6OHDA which was used in the above examples as a neurotoxin. Drug abuse with intravenous forms of drugs contaminated with MPTP, which was highly toxic to the dopaminergic neurons of the brain, has also been linked to a Parkinsonian syndrome. IGF treatment can be useful to limit brain damage in patients exposed to such neurotoxins. Environmental neurotoxins may interact with a particular genetic factor to cause the emergence of disorders or diseases such as Alzheimer's Disease.

It is appreciated that aging increases the risk of damage to the brain in many progressive neurodegenerative disorders. For example, individuals with familial forms of Alzheimer's Disease or Huntington's Disease have a genetic disorder, but the disease is not manifest typically until after the fourth or fifth decade in life. "Senile dementia" is another example. Other examples in which age is found to be a factor include but are not limited to cortical-basal ganglionic syndromes, progressive dementia, familial dementia with spastic paraparesis, progressive supranuclear palsy, and Parkinson's Disease. Age is

a risk factor in diabetic neuropathy, and this disease includes diabetic encephalopathy.

It is known that circulating IGF levels decline progressively with age (Hall and Sara (1984)). The experiments in this invention suggest that IGF levels may be progressively reduced in the brain as a consequence of this age-dependent decline in circulating IGF levels. Taken together with the knowledge that IGFs are neurotrophic factors and may be maintenance factors for the nervous system, parenteral IGF treatment is particularly helpful to effect changes in the brain, and thereby reduce the risk of damage to the brain in the various disorders in which age is a factor.

There are other brain disorders in which IGFs is expected to be useful.

In multiple sclerosis, there is a progressive demyelination of the brain and spinal cord. The etiology includes environmental factors because the prevalence is 50-fold higher in northern climes than in equatorial areas. Age plays a role, and the incidence is low in childhood and high in the third and fourth decade of life. IGFs are known to support myelination, and parenteral forms of IGFs may be useful in this disorder as well as in the diffuse cerebral sclerosis of Schilder in which demyelination is associated with progressive mental deterioration, and acute necrotizing hemorrhagic encephalomyelitis which is a fulminant form of demyelinating disease.

Various advantages and modifications will be readily apparent to those skilled in the art. The invention in its broader aspects is therefore not limited to the specific details or representative example described. Accordingly, departures may be made from the detail without departing from the spirit or scope of the disclosed general inventive concepts.

### References

U. Friedemann, "Blood-brain Barrier," *Physiol. Rev.* 22:125-45 (1942).

5 L.P. Rowland, M.E. Fink and Rubin, "Cerebrospinal Fluid: Blood-brain Barrier, Brain Edema, and Hydrocephalus," *Principles of Neural Science* (E.R. Kandel, J.H. Schwartz and T.M. Jessell eds., 3rd ed., Appleton & Lange 1991).

B. Schlosshauer, "The Blood-brain Barrier: Morphology, Molecules, and Neurothelin," *BioEssays* 15:341-46 (1993).

10 D.N. Ishii and E. Recio-Pinto, "Role of Insulin, Insulinlike Growth Factors, and Nerve Growth Factor in Neurite Formation," *Insulin, IGFs, and Their Receptors in the Central Nervous System* 315-48 (M.K. Raizada and M.A. Phillips eds., Plenum Press 1987).

15 V.R. Sara, K. Hall, H. Von Holtz, R. Humbel, B. Sjogren and L. Wetterberg, "Evidence for the Presence of Specific Receptors for Insulin-like Growth Factors 1 (IGF-1) and 2 (IGF-2) and Insulin Throughout the Adult Human Brain," *Neurosci. Lett.* 34:39-44 (1982).

C.G. Goodyear, L. De Stephano, W.H. Lai, H.J. Guyda and B.I. Posner "Characterization of Insulin-like Growth Factor Receptors in Rat Anterior Pituitary, Hypothalamus, and Brain," *Endocrinology* 114:1187-95 (1984).

20 E. Recio-Pinto, M.M. Rechler and D.N. Ishii, "Effects of Insulin, Insulinlike Growth Factor-II, and Nerve Growth Factor on Neurite Formation and Survival in Cultured Sympathetic and Sensory Neurons," *J. Neurosci.* 6:1211-19 (1986).

### References

U. Friedemann, "Blood-brain Barrier," *Physiol. Rev.* 22:125-45 (1942).

5 L.P. Rowland, M.E. Fink and Rubin, "Cerebrospinal Fluid: Blood-brain Barrier, Brain Edema, and Hydrocephalus," *Principles of Neural Science* (E.R. Kandel, J.H. Schwartz and T.M. Jessell eds., 3rd ed., Appleton & Lange 1991).

B. Schlosshauer, "The Blood-brain Barrier: Morphology, Molecules, and Neurothelin," *BioEssays* 15:341-46 (1993).

10 D.N. Ishii and E. Recio-Pinto, "Role of Insulin, Insulinlike Growth Factors, and Nerve Growth Factor in Neurite Formation," *Insulin, IGFs, and Their Receptors in the Central Nervous System* 315-48 (M.K. Raizada and M.A. Phillips eds., Plenum Press 1987).

15 V.R. Sara, K. Hall, H. Von Holtz, R. Humbel, B. Sjogren and L. Wetterberg, "Evidence for the Presence of Specific Receptors for Insulin-like Growth Factors 1 (IGF-1) and 2 (IGF-2) and Insulin Throughout the Adult Human Brain," *Neurosci. Lett.* 34:39-44 (1982).

20 C.G. Goodyear, L. De Stephano, W.H. Lai, H.J. Guyda and B.I. Posner "Characterization of Insulin-like Growth Factor Receptors in Rat Anterior Pituitary, Hypothalamus, and Brain," *Endocrinology* 114:1187-95 (1984).

25 E. Recio-Pinto, M.M. Rechler and D.N. Ishii, "Effects of Insulin, Insulinlike Growth Factor-II, and Nerve Growth Factor on Neurite Formation and Survival in Cultured Sympathetic and Sensory Neurons," *J. Neurosci.* 6:1211-19 (1986).

R. Reinhardt and C. Bondy, "Insulin Like Growth Factors Cross the Blood-brain Barrier," *Endocrinology* 135:1753-61 (1994).

P.P. Zumstein and R.E. Humbel, "Purification of Human Insulin-like Growth Factors I and II," *Methods in Enzymology* 109:782-87 (1985).

5 M.E. Svoboda and J.J. Van Wyk, "Purification of Somatomedin-C/Insulinlike Growth Factor I," *Methods in Enzymology* 109:798-816 (1985).

H.P. Guler, J. Zapf, C. Schmid and E.R. Froesch, "Insulin-like Growth Factors I and II in Healthy Man. Estimations of Half-lives and Production Rates," *Acta Endocrinologica* (Copenh) 121:753-58 (1989).

10 J. Zapf, H. Walter, E.R. Froesch, "Radioimmunological Determination of Insulin-like Growth Factors I and II in Normal Subjects and in Patients with Growth Disorders and Extrapaneatic Tumor Hypoglycemia," *J. Clin. Invest.* 68:1321-30 (1981).

G.M. Knudsen, J. Jakobsen, M. Juhler and O.B. Paulson, "Decreased  
15 Blood-brain Barrier Permeability to Sodium in Early Experimental Diabetes," *Diabetes* 35:1371-73 (1986).

J. Jakobsen, G.M. Knudsen and M. Juhler, "Cation Permeability of the Blood-brain Barrier in Streptozotocin Diabetic Rats," *Diabetologia* 30:409-13 (1987).

20 M.B. Soares, D.N. Ishii and A. Efstratiadis, "Developmental and Tissue-specific Expression of a Family of Transcripts Related to Rat Insulin-like Growth Factor II mRNA," *Nucl. Acids Res.* 13:1119-34 (1985).

D.N. Ishii, D.M. Guertin and L.R. Whalen, "Reduced Insulin-like Growth Factor-I mRNA Content in Liver, Adrenal Glands and Spinal Cord of  
25 Diabetic Rats," *Diabetologia* 37:1073-81 (1994).

M.A. Hynes, P.J. Brooks, J. Van Wyk, and P.K. Lund, "Insulin-like Growth Factor II Messenger Ribonucleic Acids are Synthesized in the Coroid Plexus of the Rat Brain," *Mol. Endocrinol.* 2:45-47 (1988).

W.H. Lee, S. Javedan and C.A. Bondy, "Coordinate Expression of Insulin-like Growth Factor System Components by Neurons and Neuroglia Cells During Retinal and Cerebellar Development," *J. Neurosci.* 12:4737-44 (1992).

5 P. Rotwein, S.K. Burgess, J.D. Milbrandt, J.E. Krause, "Differential Expression of Insulin-like Growth Factor Genes in Rat Central Nervous System," *Proc. Natl. Acad. Sci. USA* 132:167-73 (1988).

D.N. Ishii and S.B. Lupien, "Insulin-like Growth Factors Protect Against Diabetic Neuropathy: Effects on Sensory Nerve Regeneration in  
10 Rats," *J. Neurosci. Res.* 40:138-44 (1995).

H.X. Zhuang, S.F. Pu and D.N. Ishii, "Insulin-like Growth Factors (IGFs) Prevent Diabetic Neuropathy (Impaired Nerve Regeneration and Hyperalgesia) in Rats Despite Hyperglycemia," *Soc. Neurosci. Abs.* 20:415 (1994).

15 C.D. Barnes, S.J. Fung, and O. Pompeiano, "Descending Catecholaminergic Modulation of Spinal Cord Reflexes in Cat and Rat," *Ann. NY Acad. Sci.* 563:45-58 (1989).

B.E. Pulford, A.R. Mihajlov, H.O. Nornes and L.R. Whalen, "Effects of Cultured Adrenal Chromaffin Cell Implants on Catecholamine-dependent  
20 Hind Limb Reflexes in 6-OHDA Lesioned Rats," *J. Neural. Transplantation & Plasticity* 5:89-102 (1994).

A.D. Mooradian, "Diabetic Complications of the Central Nervous System," *Endocrine Reviews* 9:346-56 (1988).

A.L. McCall, "The Impact of Diabetes on the CNS," *Diabetes* 41:557-  
25 70 (1991).

E. Reske-Nielsen, K. Lundbaek, O.U. Rafaelsen, "Pathological Changes in the Central and Peripheral Nervous Systems of Young Long-term Diabetics. I. Diabetic Encephalopathy," *Diabetologia* 1:233-41 (1965).

Y. Olsson, J. Save-Soderbergh, P. Sourander and L.A. Angervall, "Patho-anatomical Study of the Central and Peripheral Nervous System in Diabetes of Early Onset and Long Duration," *Pathol. Eur.* 3:62-79 (1968).

5 H. Soininen, M. Puranen, E.L. Helkala, M. Laakson and P. Riekkinen, "Diabetes Mellitus and Brain Atrophy: A Computerized Tomography Study in an Elderly Population," *Neurobiol. Aging* 13:717-21 (1992).

10 J. Jakobsen, P. Sidenius, H.J.G. Gundersen, and R. Osterby, "Quantitative Changes of Cerebral Neocortical Structures in Insulin-treated Long-term Streptozotocin-diabetic Rats," *Diabetes* 36:597-606 (1987).

M.E. Trulson, J.H. Jacoby and R.G. MacKenzie, "Streptozotocin-induced Diabetes Reduces Brain Serotonin Synthesis in Rats," *J. Neurochem.* 46:1068-72 (1986).



## CLAIMS:

1. A method for effecting a change in the central nervous system comprising parenteral administration of an effective amount of an IGF-I.
- 5 2. A method for treating a disorder or a disease in the brain comprising parenteral administration of an effective amount of an IGF-I.
3. A method according to claim 1 or 2 wherein said IGF-I is administered  
10 in an amount from about 0.1  $\mu\text{g/kg/day}$  up to about 4  $\text{mg/kg/day}$ .
4. A method according to claim 1 or 2 wherein said IGF-I is administered in an amount from about 400  $\text{ng/kg/hour}$  up to about 160  $\mu\text{g/kg/hour}$ .
- 15 5. A method according to claim 2 wherein said disorder or disease is Alzheimer's Disease.
6. A method according to claim 2 wherein said disorder or disease is Parkinson's Disease.
- 20 7. A method according to claim 2 wherein said disorder or disease is AIDS dementia.
8. A method for effecting a change in the central nervous system comprising parenteral administration of an effective amount of an IGF-II.
- 25 9. A method for treating a disorder or a disease in the brain comprising parenteral administration of an effective amount of an IGF-II.

10. A method according to claim 8 or 9 wherein said IGF-II is administered in an amount from about 0.1  $\mu\text{g/kg/day}$  up to about 4 mg/kg/day.

5 11. A method according to claim 8 or 9 wherein said IGF-II is administered in an amount from about 400 ng/kg/hour up to about 160  $\mu\text{g/kg/hour}$ .

12. A method according to claim 9 wherein said disorder or disease is  
10 Alzheimer's Disease.

13. A method according to claim 9 wherein said disorder or disease is Parkinson's Disease.

15 14. A method according to claim 9 wherein said disorder or disease is AIDS dementia.

15. A method for effecting a change in the spinal cord comprising  
parenteral administration of an effective amount of an IGF-I or an IGF-II.

20 16. A method for treating a disorder in or trauma to the spinal cord comprising parenteral administration of an effective amount of an IGF-I or IGF-II.

25 17. A method according to claim 15 or 16 wherein said IGF-I or IGF-II is administered in an amount from about 0.1  $\mu\text{g/kg/day}$  up to about 4 mg/kg/day.

18. A method according to claim 15 or 16 wherein said IGF-I or IGF-II is administered in an amount from about 400 ng/kg/hour up to about 160  $\mu$ g/kg/hour.

1/4





Fig. 1


2/4

Fig. 2A




3/4

Fig. 2B



4/4

Fig. 2C



# INTERNATIONAL SEARCH REPORT

International Application No.  
PC1/US 96/19663

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K38/30

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ENDOCRINOLOGY, vol. 135, no. 5, November 1994, PHILADELPHIA, US, pages 1753-1761, XP000647840 R.R. REINHARDT ETAL.: "INSULIN-LIKE GROWTH FACTORS CROSS THE BLOOD-BRAIN BARRIER." cited in the application see the whole document ---	1-18
Y	WO 93 08828 A (SYNTEX-SYNERGEN NEUROSCIENCE JOINT VENTURE ET AL.) 13 May 1993 see page 8, line 11 - line 22; claims 1-3 see page 9, line 4 - line 7 see page 10, line 4 - line 13 see page 12, line 2 - line 12 see page 14, line 2 - line 10 ---	1-4,8-11
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

27 March 1997

Date of mailing of the international search report

10.04.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Ryckebosch, A



# INTERNATIONAL SEARCH REPORT

Int:      nal Application No.

PCT/US 96/19663

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 90 14838 A (CEPHALON, INC.) 13 December 1990 see page 6, line 18 - line 32 see page 8, line 26 - line 34 see page 12, line 1 - line 13 see page 20, line 5 - page 21, line 11 ---	5-7, 12-18
A	WO 93 02695 A (GENENTECH ET AL.) 18 February 1993 see claims ---	1-18
A	WO 95 13823 A (CELTRIX PHARMACEUTICALS, INC.) 26 May 1995 see page 10, line 5 - line 31; claims 1-10 -----	1-18

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 19663

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons.

1. ☒ Claims Nos.: 1-18  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim(s) 1-18  
is(are) directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.  
PCT/US 96/19663

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9308828 A	13-05-93	AU 3129793 A	07-06-93
WO 9014838 A	13-12-90	US 5093317 A	03-03-92
		CA 2058443 A	06-12-90
		EP 0476044 A	25-03-92
		JP 7068138 B	26-07-95
		JP 4507240 T	17-12-92
WO 9302695 A	18-02-93	CA 2114251 A	18-02-93
		EP 0597033 A	18-05-94
WO 9513823 A	26-05-95	AU 1057095 A	06-06-95
		CA 2176708 A	26-05-95
		EP 0729361 A	04-09-96